

EXISTENCE AND REGULARITY OF MONOTONE SOLUTIONS TO A FREE BOUNDARY PROBLEM

DANIELA DE SILVA

ABSTRACT. We investigate existence and regularity properties of one-phase free boundary graphs, in connection with the question of whether there exists a complete non-planar free boundary graph in high dimensions.

1. INTRODUCTION

Let Ω be an open connected subset of $\mathbb{R}^n$, and consider the energy functional

$$J(u, \Omega) = \int_{\Omega} (|\nabla u|^2 + \chi_{\{u>0\}}).$$

In [AC], Alt and Caffarelli analyzed the question of the existence and regularity of a minimizer u of $J(\cdot, \Omega)$. They developed a partial regularity theory for the free boundary of u , that is

$$F(u) = \partial\{u > 0\} \cap \Omega,$$

showing that its reduced part $F^*(u)$ is locally $C^{1,\alpha}$. Higher regularity results of Kinderlehrer and Nirenberg [KN], then imply that $F^*(u)$ is locally analytic. In [AC] the authors also proved that in two dimensions, $F(u)$ does not have singularities. Subsequently in [W2], Weiss showed that there exists a critical dimension k , $3 \leq k \leq +\infty$, such that energy minimizing free boundaries are smooth for $n < k$. These results draw on a strong analogy with the theory of minimal surfaces, for which it is known that the critical dimension is 8. In [CJK], Caffarelli, Jerison and Kenig proved that there are no singular free boundary minimizers in dimension $n = 3$, which yields $k \geq 4$. Their proof suggests that $k = 7$, but the problem remains still open. In [DJ] De Silva and Jerison showed that $k \leq 7$, by providing the first example of a singular energy minimizing free boundary in dimension $n = 7$. Analogously, for the theory of minimal surfaces, the Simons cone, provides an example of a singular set of minimal perimeter in dimension $n = 8$.

The purpose of this note is to pursue even further the analogy between the theory of minimal surfaces and free boundary regularity, precisely we turn to free boundary graphs. Our motivation lies in the question of whether there exists a complete non-planar free boundary graph, i.e a classical solution on $\mathbb{R}^n$ to the problem,

$$(1.1) \quad \begin{cases} \Delta u = 0 & \text{in } \{u > 0\}, \\ |\nabla u| = 1 & \text{on } \partial\{u > 0\}, \\ \partial\{u > 0\} & \text{is a non-planar graph in the } x_n \text{ direction.} \end{cases}$$

This is the analogue of the celebrated Bernstein problem for minimal graphs. Precisely, it is known that planes are the only complete smooth minimal graphs in $\mathbb{R}^n$,

when $n \leq 8$ (see for example [G]). This result is sharp since in [BDG], Bombieri, De Giorgi and Giusti proved the existence of a non affine minimal graph in dimension 9, which turns out to be strictly related to the existence of the Simons cone, one dimension lower. The result in [DJ], then naturally raises the analogous question for free boundary graphs. In analogy with the minimal surfaces theory, we expect that a global smooth solution to (1.1) exists in dimension 8 or higher.

The first step towards constructing such an example, is to develop a local theory which is the analogue of the existence and regularity theory for the minimal surface equation in a ball, when the boundary data is smooth. Then, a limiting argument provides a global solution. In order to prevent this global solution from being planar, one wishes to control its behavior by trapping it between a given global subsolution and a given global supersolution. However, since ordinary (strong) comparison results are not available in the free boundary context (see Lemma 4.4), that is if two solutions are one greater than the other one on the boundary then the inequality may not be preserved in the interior, this “trapping” is not straightforward. Thus, we develop our local theory by constructing a solution which is trapped in between a given subsolution and supersolution. Moreover, since we wish to preserve the graph property in the limit, we also need to construct a local solution which enjoys a certain regularity property (density property), which for example, would not be guaranteed if we were to construct our solution via a standard Perron method. For this reason, we seek a solution which is also an energy minimizer to the energy functional J among a certain class of competitors.

Our result is the following. Let $R, h_R > 0$, and let C_R denote the cylinder $C_R = \mathcal{B}_R(0) \times \{|x_n| < h_R\}$, and S_R denotes the sides of the cylinder C_R , $S_R = \partial\mathcal{B}_R(0) \times \{|x_n| \leq h_R\}$; consider the one-phase free boundary problem,

$$(1.2) \quad \begin{cases} \Delta u = 0 & \text{in } C_R^+(u) := \{x \in C_R : u(x) > 0\}, \\ |\nabla u| = 1 & \text{on } F(u) := (\partial C_R^+(u)) \cap C_R. \end{cases}$$

Theorem 1.1. *Assume that, there exist a strict smooth subsolution V_1 and a strict smooth supersolution V_2 to (1.1) in $\mathbb{R}^n$, such that*

- (i) $0 \leq V_1 \leq V_2$ on $\mathbb{R}^n$; $0 \in \{V_2 > 0\} \cap \{V_1 = 0\}^\circ$;
- (ii) $\partial_n V_i > 0$ in $\overline{\{V_i > 0\}}$, for $i = 1, 2$.

Then, for each $R > 0$ and h_R sufficiently large, there exists u_R viscosity solution to (1.2), such that u_R is minimizes J over $K_R := \{v \in H^1(C_R) \mid V_1 \leq v \leq V_2 \text{ a.e on } C_R, v = V_2 \text{ on } S_R\}$, u_R is monotone increasing in $C_R^+(u)$ in the x_n direction, and $V_1 \leq u_R \leq V_2$. Moreover, in the interior, $F(u_R)$ is a smooth graph in the x_n direction.

For the precise definition of viscosity solution, we refer the reader to Section 2. We remark that the proof of the result in [DJ], provides a clear indication of how to construct functions V_1 and V_2 satisfying the assumptions above, when $n \geq 8$. We plan on constructing these functions in a future paper.

The main tools to achieve the existence part in our Theorem are blow-up and domain variation techniques. Then, the fact that u_R is also a viscosity solution, allows us to use maximum principle techniques and a continuity argument, to compare u_R with a family of subsolutions, which are suprema of vertical translates of

u_R over balls (supconvolutions). This yields the desired Lipschitz behavior (hence smoothness) of the free boundary of u_R .

As already observed, the second step towards constructing a global solution to (1.1) is a limiting argument as $R \rightarrow +\infty$. In the theory of minimal surfaces, the convergence to a global solution is guaranteed by a very powerful tool, that is the a-priori estimate of the gradient of a solution to the minimal surface equation. In a forthcoming paper, we prove the analogue of such a tool in the free boundary context [DJ2].

Here, in order to preserve the graph property in the limit, we prove that the positive phase of our solution is a non-tangentially accessible (NTA) domain, that is, it enjoys a certain scale-invariant connectivity property (see Section 2 for the precise definition of NTA domains.) Then, a limiting argument allows us to prove the following:

Theorem 1.2. *Assume that, there exist a strict smooth subsolution V_1 and a strict smooth supersolution V_2 to (1.1) in $\mathbb{R}^n$, such that:*

- (i) $0 \leq V_1 \leq V_2$ on $\mathbb{R}^n$;
- (ii) $\partial_n V_i > 0$ on $\{V_i > 0\}$, $i = 1, 2$;
- (iii) $\lim_{r \rightarrow \infty} \frac{V_1(rx)}{r} \geq U(x')$.

Then, there exists a global energy minimizing viscosity solution u to:

$$\Delta u = 0 \quad \text{in} \quad \{u > 0\}, \quad |\nabla u| = 1 \quad \text{on} \quad F(u),$$

such that u is monotone increasing in $\overline{\{u > 0\}}$ in the x_n direction, and $F(u)$ is a continuous non-planar graph, with a universal modulus of continuity on each compact subset of $\mathbb{R}^n$. Moreover, $F(u)$ is NTA.

Here U is the singular global minimizer in $\mathbb{R}^{n-1}$ from [DJ], interpreted as a function of n variables. Hypothesis (iii) is used only to prevent $F(u)$ from being planar. While we could weaken this assumption, its motivation lies in the fact that, in analogy with the minimal surfaces theory, we expect a smooth non-affine free boundary graph u to blow down to an energy minimizing solution.

The NTA property of $F(u)$ is proved by the means of a monotonicity formula [ACF] for ∇u , together with non-degeneracy properties of u . The proof follows from arguments in [ACS]. Then, exploiting the known behavior of positive harmonic functions in NTA domains [JK], we derive that $F(u)$ cannot contain vertical segments.

The paper is organized as follows. In Section 2, we introduce some notation and definitions. In Section 3, we show the existence of a “trapped” monotone energy minimizing viscosity solution to our one-phase free boundary problem on a cylinder. Then, in Section 4, we show that such a solution is indeed smooth in the interior. In Section 5, we prove that the free boundary of our solution is locally NTA. Finally, in Section 6, we prove the existence of a global monotone viscosity solution whose free boundary is a graph in the vertical direction, with a universal modulus of continuity on each compact of $\mathbb{R}^n$, trapped between two given graphs.

2. NOTATION AND DEFINITIONS.

In this section we collect some notation and definitions which will be used throughout this paper.

A point $x \in \mathbb{R}^n$ will be denoted by (x', x_n) , with $x' = (x_1, \dots, x_{n-1})$. A ball of radius r in $\mathbb{R}^{n-1}$, will be denoted by $\mathcal{B}_r$, while a ball of radius r in $\mathbb{R}^n$, will be denoted by B_r . When specifying the center x of the ball, we will use either $B_r(x)$ or $B(x, r)$. Also, Ω denotes an open bounded connected subset of $\mathbb{R}^n$.

For any non-negative function u on Ω , set

$$\Omega^+(u) = \{x \in \Omega : u(x) > 0\}, \quad \Omega^- = \{x \in \Omega : u = 0\}^\circ, \quad F(u) = (\partial\Omega^+(u)) \cap \Omega.$$

Consider the one-phase free-boundary problem:

$$(2.1) \quad \begin{cases} \Delta u = 0 & \text{in } \Omega^+(u), \\ |\nabla u| = 1 & \text{on } F(u). \end{cases}$$

We recall the following standard definition (see for example [C1].)

Definition 2.1. Let v be a nonnegative continuous function in Ω . We say that v is a viscosity subsolution (resp. supersolution) to (2.1) in Ω , if and only if the following conditions are satisfied:

- (i) $\Delta v \geq 0$ (resp. ≤ 0) in $\Omega^+(v)$;
- (ii) If $x_0 \in F(v)$ and $F(v)$ has at x_0 a tangent ball B_ϵ from the positive (resp. zero) side (i.e. $B_\epsilon \subset \Omega^+(v)$ (resp. $\Omega^-(v)$), $x_0 \in \partial B_\epsilon$), then, for some $\alpha \geq 1$ (resp. $\alpha \leq 1$) and ν the unit inner (resp. outer) radial direction of ∂B_ϵ at x_0 ,

$$v(x) = \alpha(x - x_0, \nu)^+ + o(|x - x_0|), \text{ as } x \rightarrow x_0.$$

If the constant α in Definition 2.1 is strictly greater (resp. smaller) than 1, then u is called a strict viscosity subsolution (resp. supersolution).

When u is simultaneously a viscosity subsolution and a viscosity supersolution, then u is called a viscosity solution.

Our viscosity solution will satisfy certain regularity properties, which we now define. Let u be a continuous non-negative function on Ω . Set,

$$d(x) = \text{dist}(x, F(u)).$$

Definition 2.2. We say that u is non-degenerate, if and only if, for every $G \Subset \Omega$, there exists a constant $K = K(G)$ such that

$$u(x) \geq Kd(x),$$

for all $x \in G^+(u)$, with $B_{d(x)}(x) \subset G$.

Definition 2.3. We say that u is (I) non-degenerate, if and only if, for every $G \Subset \Omega$, there exists a constant $K = K(G)$ such that, for any ball $B_r \subset G$ centered at a free boundary point,

$$\int_{B_r} u \geq Kr.$$

Definition 2.4. We say that $F(u)$ satisfies the density property (D) if and only if:

- (D) for any $G \Subset \Omega$, there exists a constant $c = c(G) < 1$, such that, for any ball $B_r \subset G$ centered at a free boundary point,

$$c \leq \frac{|B_r \cap \{u > 0\}|}{|B_r|} \leq 1 - c.$$

Such properties are crucial to use so-called “blow-up” techniques. Specifically, let u be a non-negative, Lipschitz continuous function in Ω . Let $x_0 \in F(u)$, and let $B_{r_k}(x_0) \subset \Omega$ be a sequence of balls with $r_k \rightarrow 0$, as $k \rightarrow +\infty$. Consider the blow-up sequence:

$$u_k(x) = \frac{1}{r_k} u(x_0 + r_k x).$$

Since for a given $D \Subset \mathbb{R}^n$ and large k the functions u_k are uniformly Lipschitz continuous in D , there exists a function $u_0 : \mathbb{R}^n \rightarrow \mathbb{R}$, such that:

$$\begin{aligned} u_k &\rightarrow u_0 \quad \text{in } C_{loc}^{0,\alpha}(\mathbb{R}^n), \quad \text{for all } 0 < \alpha < 1; \\ \nabla u_k &\rightarrow \nabla u_0 \quad \text{weakly star in } L_{loc}^\infty(\mathbb{R}^n). \end{aligned}$$

Moreover, u_0 is Lipschitz continuous in the entire space. The globally defined function u_0 is called a blow-up of u . Using the same argument as in [F], (see Chapter 3, Lemma 3.6), one can prove the following.

Lemma 2.5. *Let u be a non-negative function in Ω , harmonic in $\Omega^+(u)$, Lipschitz continuous and (I) non-degenerate. Assume that u satisfies the density property (D). Then the following properties hold:*

- (a) $\partial\{u_k > 0\} \rightarrow \partial\{u_0 > 0\}$ in the Hausdorff distance;
- (b) $\chi_{\{u_k > 0\}} \rightarrow \chi_{\{u_0 > 0\}}$ in L_{loc}^1 ;
- (c) $\nabla u_k \rightarrow \nabla u_0$ a.e.

Moreover, u_0 is (I) non degenerate, and it satisfies the density property (D).

We also need the following result from [W1], which characterizes blow-up limits of certain solutions (variational solutions) to the free boundary problem (2.1).

First we recall the following definition. Let,

$$(2.2) \quad J(u, \Omega) = \int_{\Omega} (|\nabla u|^2 + \chi_{\{u > 0\}}).$$

Definition 2.6. We define $u \in H_{loc}^1(\Omega)$ to be a variational solution to (2.1), if $u \in C(\Omega) \cap C^2(\Omega^+(u))$ and

$$0 = -\frac{d}{d\epsilon} J(u(x + \epsilon \eta(x)))|_{\epsilon=0} = \int_{\Omega} (|\nabla u|^2 \operatorname{div} \eta - 2 \nabla u D \eta \nabla u + \chi_{\{u > 0\}} \operatorname{div} \eta)$$

for any $\eta \in C_0^1(\Omega, \mathbb{R}^n)$.

Lemma 2.7. *Let u be a variational solution to (2.1) in Ω , and assume that u is Lipschitz continuous and satisfies the density property (D). Then any blow up limit of u is homogeneous of degree 1.*

Such blow-up technique will allow us to prove that our energy minimizing solution is indeed a trapped viscosity solution, which also enjoys the desired regularity properties. These regularity properties in Definitions (2.2)-(2.3)-(2.4), are crucial in the proof that the free boundary of our solution is locally NTA.

We conclude this section by recalling the notion of Non-Tangentially Accessible (NTA) domains (see [JK]). Let D be a bounded domain in $\mathbb{R}^n$. A M -non-tangential ball in D , is a ball $B_r \subset D$, such that: $Mr > \operatorname{dist}(B_r, \partial D) > M^{-1}r$. For $x_1, x_2 \in D$, a Harnack chain from x_1 to x_2 in D is a sequence of M -non-tangential balls, such that the first ball contains x_1 , the last contains x_2 , and such that consecutive balls intersect.

Definition 2.8. A bounded domain D in $\mathbb{R}^n$ is called NTA, when there exist constants M and $r_0 > 0$ such that:

- (i) Corkscrew condition. For any $x \in \partial D$, $r < r_0$, there exists $y = y_r(x) \in D$ such that $M^{-1}r < |y - x| < r$ and $\text{dist}(y, \partial D) > M^{-1}r$;
- (ii) The Lebesgue density of D^c at any of its points is bounded below uniformly by a positive constant C , i.e for all $x \in \partial D$, $0 < r < r_0$,

$$\frac{|B(x, r) \setminus D|}{|B(x, r)|} \geq C;$$

- (iii) Harnack chain condition. If $\epsilon > 0$ and x_1, x_2 belong to D , $\text{dist}(x_j, \partial D) > \epsilon$ and $|x_1 - x_2| < C\epsilon$, then there exists a Harnack chain from x_1 to x_2 whose length depends on C , but not on ϵ .

3. LOCAL EXISTENCE THEORY.

In this section we prove the existence of an energy minimizing viscosity solution u_R as in Theorem 1.1. We also show that u_R enjoys the regularity properties from Definitions (2.2)-(2.3)-(2.4). The regularity of the free boundary $F(u_R)$ will be investigated in the next section.

Precisely, for $R, h_R > 0$, let

$$C_R := \mathcal{B}_R(0) \times \{|x_n| < h_R\}, \quad S_R = \partial \mathcal{B}_R(0) \times \{|x_n| \leq h_R\}.$$

We prove the following result.

Theorem 3.1. *Assume that, there exist a strict smooth subsolution V_1 and a strict smooth supersolution V_2 to (2.1) in $\mathbb{R}^n$, such that*

- (i) $V_1 \leq V_2$ on $\mathbb{R}^n$, $0 \in \{V_2 > 0\} \cap \{V_1 = 0\}^\circ$;
- (ii) $\partial_n V_i > 0$ in $\overline{\{V_i > 0\}}$, for $i = 1, 2$.

Then, for each $R > 0$, and h_R sufficiently large, there exists u_R minimizer of J over $K_R := \{v \in H^1(C_R) | V_1 \leq v \leq V_2 \text{ a.e on } C_R, v = V_2 \text{ on } S_R\}$, such that

- (a) u_R is monotone increasing in $C_R^+(u)$ in the x_n direction;
- (b) u_R is Lipschitz continuous, non-degenerate, (I) non-degenerate, and satisfies the density property (D) in C_R ;
- (c) u_R is a viscosity solution to (2.1) in C_R .

Remark 1. The height h_R of the cylinder C_R , must be such that the free boundaries of V_1 and V_2 “exit” from the sides of the cylinder. For example, let $\partial\{V_i > 0\} = \{(x', \phi_i(x')), x' \in \mathbb{R}^{n-1}\}$. Then we can choose

$$h_R \geq \max\{2d_R(V_1), 2d_R(V_2), R\}, \quad d_R(V_i) = \max_{\mathcal{B}_R(0)} |\phi_i(x')|, i = 1, 2.$$

Remark 2. Under the assumptions of Theorem 3.1, $V_1 < V_2$ on $\{V_2 > 0\}$, and $F(V_1) \cap F(V_2) = \emptyset$.

PROOF. Assume that there exists $x \in \{V_2 > 0\}$ such that, $V_1(x) = V_2(x)$. Then, since V_1 is subharmonic in $\{V_2 > 0\}$, the maximum principle implies $V_1 \equiv V_2$, which contradicts the fact that V_1 is a strict subsolution and V_2 is a strict supersolution. Analogously, suppose $x \in F(V_1) \cap F(V_2)$, and let $B \subset \{V_1 > 0\}$ be a ball tangent to $F(V_1)$ at x . Then, by Hopf’s lemma, $\partial_\nu(V_1 - V_2) < 0$, with ν inner normal derivative

to ∂B at x . Again, this contradicts the fact that V_1 is a strict subsolution, and V_2 is a strict supersolution. $\square$

Remark 3. One could prove that if $u \geq 0$ is a Lipschitz continuous function on Ω , and u is harmonic on $\Omega^+(u)$, then the two notions of non-degeneracy given in Definition 2.2 and Definition 2.3 are equivalent (see also Lemma 7 in [C3]). We do not prove this fact here, since according to part (b) in Theorem 3.1, we can prove directly that u_R satisfies both properties. Therefore, we think of these two definitions as two versions of the same property.

Let $W_R := C_R^+(V_2) \cap C_R^-(V_1)$ be the region between the two smooth (non-touching) graphs $F(V_1), F(V_2)$. Observe that if u_R minimizes J over K_R , then it minimizes $J(\cdot, W_R)$ among all competitors which equal u_R on ∂W_R . Hence, u_R is Lipschitz continuous, non-degenerate and satisfies the density property (D) in W_R (see [AC]). Thus, in the proof of part (b) in Theorem 3.1, we only need to analyze what happens when $F(u_R)$ is “close” to the fixed boundary of W_R , that is either to $F(V_1)$ or to $F(V_2)$. Whenever we are “away” from either $F(V_1)$ or $F(V_2)$ the desired properties will follow with the same arguments as in [AC] or [ACF], while near $F(V_1)$ or $F(V_2)$, the free boundary $F(u_R)$ will inherit the same good properties of the smooth graphs $F(V_1), F(V_2)$.

Moreover, according to Lemma 7 in [C2], u_R is also a viscosity solution to (2.1) in W_R . Thus, in order to show part (c) in Theorem 3.1 it is enough to show that in fact $F(u_R) \subset W_R$, that is $F(u_R)$ does not touch neither $F(V_1)$ nor $F(V_2)$.

PROOF OF THEOREM 3.1. We divide the proof in four steps.

Step 1: Existence of monotone minimizers. We recall that for any Lebesgue measurable function u on $\overline{C_R}$ one can define the monotone (non-decreasing) rearrangement u^* of u , in the direction x_n . The function u^* is monotone non-decreasing in the x_n -direction, and u and u^* are equimeasurable, that is, for all $k \in \mathbb{R}$

$$|\{u \geq k\}| = |\{u^* \geq k\}|.$$

Moreover, the mapping $u \rightarrow u^*$ is order preserving, i.e, $u \leq v$ implies $u^* \leq v^*$. The following proposition holds (for details we refer the reader to [K]).

Proposition 3.2. *Let $u \in W^{1,p}(C_R)$, $1 < p < \infty$. Then $u^* \in W^{1,p}(C_R)$ and we have*

$$(3.1) \quad \int_{C_R} |\nabla u|^p dx \geq \int_{C_R} |\nabla u^*|^p dx.$$

Now, consider the energy functional $J_R(\cdot) = J(\cdot, C_R)$ defined in (2.2). Denote by K_R the following closed and convex subset of $H^1(C_R)$,

$$K_R := \{v \in H^1(C_R) | V_1 \leq v \leq V_2 \text{ a.e on } C_R, v = V_2 \text{ on } S_R\}.$$

The following existence theorem holds.

Theorem 3.3. *The minimum of J_R over K_R is achieved at u_R , which is a monotone non-decreasing function in $C_R^+(u)$ in the x_n -direction.*

PROOF. Since J_R is non-negative, there exists a minimizing sequence u_m , that is

$$u_m \in K_R, \quad J_R(u_m) \rightarrow \alpha \equiv \inf_{v \in K_R} J_R(v), \quad 0 \leq \alpha \leq J_R(V_2) < \infty.$$

The sequence $\{u_m\}$ is uniformly bounded in $H^1(C_R)$. Indeed,

$$\|\nabla u_m\|_2^2 \leq J(V_2), \quad \|u_m\|_2 \leq \|V_2\|_2.$$

Therefore, we can extract a subsequence, which we will still denote by $\{u_m\}$, such that $u_m \rightarrow u \in K_R$, weakly in $H^1(C_R)$. It is known that J_R is lower semicontinuous (see [AC]) with respect to weak H^1 convergence, that is,

$$\liminf_{m \rightarrow \infty} J_R(u_m) \geq J_R(u).$$

This immediately implies that u is a minimizer for J_R over K_R . Now, let u^* be the monotone rearrangement of u . Then, using Proposition 3.2, together with the equimeasurability of rearrangements, we get that

$$J_R(u^*) \leq J_R(u).$$

Moreover, the order preserving property implies that $u^* \in K_R$. Hence u^* is the desired minimizer, monotone non-decreasing in the x_n direction. $\square$

We will henceforth denote by u_R , a minimizer of J_R over K_R , which is monotone non-decreasing in the x_n direction.

Step 2: Continuity and harmonicity of monotone minimizers. We wish to show that u_R is harmonic in its positive phase. This is achieved via standard techniques.

Lemma 3.4. *u_R is continuous in $C_R^+(V_2)$, and harmonic in $C_R^+(u_R)$. In particular, u_R is monotone increasing in $C_R^+(u_R)$ in the x_n direction.*

PROOF. Let D be a compact subset of $C_R^+(V_2)$, and let B_ρ be a ball of radius ρ in D . Denote by v_ρ the harmonic replacement of u_R on B_ρ , that is the harmonic function in B_ρ which equals u_R on ∂B_ρ . Assume that v_ρ is extended to be u_R outside B_ρ . Since $0 \leq u_R \leq V_2$ a.e., we have $0 \leq (v_\rho - V_2)^+ \leq (v_\rho - u_R)^+$. Hence $(v_\rho - V_2)^+ \in H_0^1(B_\rho)$ (see [AH]). Therefore, by the weak maximum principle (see [GT]) we obtain $v_\rho \leq V_2$ a.e. on B_ρ . Analogously, we get $V_1 \leq v_\rho$ a.e. on B_ρ . Since u_R minimizes $J(\cdot, B_\rho)$ among all competitors v , such that $V_1 \leq v \leq V_2$, and $v = u_R$ on ∂B_ρ , we get that

$$\int_{B_\rho} (|\nabla u_R|^2 + \chi_{\{u_R > 0\}}) \leq \int_{B_\rho} (|\nabla v_\rho|^2 + \chi_{\{v_\rho > 0\}}).$$

Therefore,

$$\int_{B_\rho} (|\nabla u_R|^2 - |\nabla v_\rho|^2) \leq K\rho^n.$$

A standard iterative argument (see [ACF]) then implies the desired continuity. Now, take $\bar{x} \in C_R^+(u_R)$. By continuity, there exists $r > 0$ such that $B_r(\bar{x}) \subset C_R^+(u_R)$. Let w_r , be the harmonic replacement of u_R on $B_r(\bar{x})$. Since w_r minimizes the Dirichlet integral and $w_r > 0$ on $B_r(\bar{x})$, we get that

$$J(w_r, B_r(\bar{x})) \leq J(u_R, B_r(\bar{x})).$$

As before, the minimality of u_R implies that the reverse inequality holds as well. Hence

$$\int_{B_r(\bar{x})} |\nabla w_r|^2 = \int_{B_r(\bar{x})} |\nabla u_R|^2.$$

By uniqueness of the Dirichlet minimizer we obtain then $u_R = w_r$ on $B_r(\bar{x})$. $\square$

Step 3: Lipschitz continuity, non-degeneracy, and density property of monotone minimizers. We start by proving the Lipschitz continuity of u_R in C_R . Set $d(x, K) = \text{dist}(x, K)$, for any closed set K . In particular,

$$d(x) = d(x, F(u_R)).$$

Lemma 3.5. u_R is Lipschitz continuous in C_R , with universal constant on each $D \Subset C_R$. In particular, for every $D \Subset D' \Subset C_R$, there exists $K > 0$ depending on D, n and on the C^1 norm of V_2 on D' , such that,

$$u_R(x) \leq Kd(x), \quad \text{for all } x \in D.$$

PROOF. Let $x_0 \in D \Subset D' \Subset C_R$, with $u(x_0) > 0$, and let $B_r = B_r(x_0)$ be the maximum ball contained in $D' \cap \{u > 0\}$. If ∂B_r touches $\partial D'$, then $r \geq \text{dist}(D, D')$, and we can apply interior regularity together with the fact that $u_R \leq V_2$, in order to show $|\nabla u_R|(x_0) \leq K$. Otherwise, ∂B_r touches $F(u_R)$ at a point x_1 .

We distinguish two cases.

(a) $d(x_1, F(V_2)) > r/2$. Then $B_{r/2}(x_1) \subset C_R^+(V_2)$ and we can apply the argument of Lemma 3.2 from [AC] to conclude the following,

$$\int_{B_{r/2}(x_1)} u_R dx \leq Kr.$$

Now, let $\bar{x}$ be on the ray from x_0 to x_1 , at distance $r/4$ from x_1 . Then, by Harnack inequality, and the mean value property for u_R , we get (with K changing for each inequality)

$$u_R(x_0) \leq Ku_R(\bar{x}) = K \int_{B_{r/4}(\bar{x})} u_R \leq K \int_{B_{r/2}(x_1)} u_R \leq Kr.$$

(b) $d(x_1, F(V_2)) \leq r/2$. Then,

$$u_R(x_0) \leq V_2(x_0) \leq Kd(x_0, F(V_2)) \leq K|x_0 - x_1| + d(x_1, F(V_2)) \leq Kr,$$

where K depends on the Lipschitz norm of V_2 on D' .

Now, denote by $v(x) = u_R(rx + x_0)/r$. Then, $\Delta v = 0$ and by Harnack inequality, $v(x) \leq K$ on $B_{1/2}(0)$. By interior regularity, $|\nabla v| \leq K'$ on $B_{1/4}(0)$, with K' dimensional constant. Rescaling back to u_R , we obtain $|\nabla u_R| \leq K'$, on $B_{r/4}(x_0)$, which implies the desired Lipschitz continuity. $\square$

Corollary 3.6. u_R is a Lipschitz continuous subharmonic function in C_R .

We now prove a non-degeneracy result.

Lemma 3.7. u_R is non-degenerate on C_R , i.e., for every $D \Subset D' \Subset C_R$, there exists $\bar{K} > 0$ depending on D, n and on the C^2 norm of V_1 on D' , such that

$$\bar{K}d(x) \leq u_R(x),$$

for all $x \in D^+(u_R)$, such that $B_{d(x)}(x) \subset D$.

PROOF. Let $x_0 \in D^+(u_R)$, and denote by $r = d(x_0)$. Assume $B_r(x_0) \subset D$. We distinguish two cases.

(a) $d(x_0, F(V_1)) > r/2$. If $x_0 \in C_R^-(V_1)$, then $B_{r/2}(x_0) \subset C_R^-(V_1)$ and we can proceed as in [ACF] Theorem 3.1, to conclude that

$$u_R(x_0) = \int_{B_{r/2}(x_0)} u_R \geq \bar{K}r.$$

If $V_1(x_0) > 0$, then

$$u_R(x_0) \geq V_1(x_0) \geq Kd(x_0, F(V_1)) \geq \bar{K}r,$$

with K depending on the C^2 norm of V_1 on D' .

(b) $d(x_0, F(V_1)) = |x_0 - x_1| \leq r/2$. Then $B_{r/2}(x_1) \subset B_r(x_0)$, hence by Harnack inequality,

$$u_R(x_0) \geq K \sup_{B_{r/4}(x_1)} u_R \geq K \sup_{B_{r/4}(x_1)} V_1 \geq \bar{K}r,$$

again with $\bar{K}$ depending on the C^2 norm of V_1 on D' . $\square$

We wish to prove a density property for free boundary points. Towards this aim, we will need to reformulate our non-degeneracy property in the following way:

Corollary 3.8. *u_R is (I) non-degenerate on C_R , i.e. for any $D \Subset C_R$, there exist a constant K , such that, for any ball $B_r \subset D$ centered at a free boundary point,*

$$\int_{B_r} u_R \geq Kr.$$

The corollary above can be deduced by the arguments in the proof of Lemma 3.7. We are now ready to derive the following density property.

Lemma 3.9. *u_R satisfies the density property (D) on C_R , i.e. for any $G \Subset G' \Subset C_R$, there exist a constant $c < 1$, depending on G, n and on the Lipschitz constant of $F(V_2)$ on G' , such that for any ball $B_r \subset G$ centered at a free boundary point,*

$$c \leq \frac{|B_r \cap \{u_R > 0\}|}{|B_r|} \leq 1 - c.$$

PROOF. Assume B_r is centered at 0. By Corollary 3.8, there exists $y \in \partial B_{r/2}$ such that, $u(y) \geq Kr/2$. By Lipschitz continuity, for any $z \in B_{kr}(y)$ we have:

$$u_R(z) \geq u_R(y) - C|z - y| > Kr/2 - Ckr > 0$$

as long as k is sufficiently small. Hence $B_{kr}(y) \subset B_r \cap \{u_R > 0\}$, from which the desired lower bound follows. In order to get the upper bound, we distinguish two cases.

(a) $d(0, F(V_2)) = |x_0| \leq r/2$. Then $B_{r/2}(x_0) \subset B_r$. Hence,

$$|\{u = 0\}^\circ \cap B_r| \geq |\{V_2 = 0\}^\circ \cap B_{r/2}(x_0)| \geq C|B_{r/2}(x_0)| \geq C'|B_r|.$$

(b) $d(0, F(V_2)) = |x_0| > r/2$. Then, $B_{r/2}(0) \subset \{V_2 > 0\}$. Hence we can replace u_R with its harmonic replacement on $B_{r/2}(0)$, and proceed as in [AC], Lemma 3.7. $\square$

Remark. One could prove the statements of Lemma 3.5 and Lemma 3.7 with the constants $K, \bar{K}$ independent of V_2 , and V_1 respectively.

Lemmas 3.5-3.7-3.9 and Corollary 3.8 prove the statement in part (b) of our Theorem.

Step 4: Non-touching of the free boundaries. We are now ready to prove the following statement, using blowing-up techniques.

Lemma 3.10. $F(u_R)$ does not intersect neither $F(V_1)$ nor $F(V_2)$.

We start by proving that u_R is separated from V_1 and V_2 in its positive phase.

Lemma 3.11. $u_R < V_2$ in $C_R^+(V_2)$, and $V_1 < u_R$ in $C_R^+(u_R)$.

PROOF. Assume $u_R(x) = V_2(x)$ at some point $x \in C_R^+(V_2)$, then the strong maximum principle implies that $u_R \equiv V_2$ on $C_R^+(V_2)$, hence $u_R \equiv V_2$ on C_R . We want to show that this contradicts the fact that u_R minimizes J_R on K_R . Let $g \in C_0^\infty(C_R)$, $g \leq 0$. For $\epsilon > 0$, set $y_\epsilon(x) = x + \epsilon g e_n$ and $V_\epsilon(x) = u_R(y_\epsilon(x))$. For ϵ sufficiently small, the monotonicity of V_2 in the x_n -direction and the fact that $V_1 < V_2$ in the positive phase of V_2 , imply that $V_\epsilon \in K_R$. Therefore, using that $\text{Det}(y_\epsilon(x)) = 1 + \epsilon \nabla \cdot g e_n + o(\epsilon^2)$, we get that

$$\begin{aligned} 0 &\leq J_R(V_\epsilon) - J_R(u_R) = \\ &= \epsilon \left\{ \int_{C_R} -(|\nabla u_R|^2 + \chi_{\{u_R > 0\}}) \nabla \cdot g e_n + (2 \nabla u_R D g e_n \nabla u_R) \right\} + o(\epsilon^2). \end{aligned}$$

Therefore using Lemma 3.4, we obtain ($u_R \equiv V_2$ and V_2 smooth)

$$\begin{aligned} 0 &\geq \int_{\{V_2 > 0\}} \nabla \cdot ((|\nabla V_2|^2 + 1) g e_n - 2 g e_n \cdot \nabla V_2 \nabla V_2) = \\ &= - \int_{\partial\{V_2 > 0\}} ((|\nabla V_2|^2 + 1) g e_n - 2 g e_n \cdot \nabla V_2 \nabla V_2) \cdot \nu = \\ &= - \int_{\partial\{V_2 > 0\}} (1 - |\nabla V_2|^2) g \nu_n \end{aligned}$$

for all function g as above, and ν the inner unit normal to $\partial\{V_2 > 0\}$. This contradicts the strict supersolution property of V_2 .

Assuming now, $u_R(x) = V_1(x)$ at some point $x \in C_R^+(u_R)$, then the contradiction follows immediately by the fact that $V_1 < V_2$ on $\{V_2 > 0\}$, and $u_R = V_2$ on S_R . $\square$

PROOF OF LEMMA 3.10. We prove that $F(u_R)$ does not touch $F(V_2)$. The proof that $F(u_R)$ does not intersect $F(V_1)$ follows by similar arguments.

Assume by contradiction that there exists $x_0 \in F(u_R) \cap F(V_2)$. Then, $F(u_R)$ has a tangent ball from the zero side at x_0 , and according to Lemma A1 in [C2], $u_R(x) = a(x - x_0, \nu)^+ + o(|x - x_0|)$, near x_0 , from the positive side of u_R , with ν the inner normal to $F(V_2)$ at x_0 . Furthermore, by non-degeneracy (Lemma 3.7), $a > 0$. Let $B_{\rho_k}(x_0)$ be a sequence of balls with $\rho_k \rightarrow 0$ such that $u_k(x) := \frac{1}{\rho_k} u_R(x_0 + \rho_k x)$ blows up to $U(x)$, and $V_k(x) := \frac{1}{\rho_k} V_2(x_0 + \rho_k x)$ blows up to $\overline{V_2}$. Thus, on the unit ball B , $U(x) = a(x, \nu)^+$ and $\overline{V_2}(x) = b(x, \nu)^+$, and

$$(3.2) \quad 0 < a \leq b < 1.$$

Let $B_\delta(x_0)$ be a ball centered at x_0 , with δ small enough so that $B_\delta(x_0) \subset C_R^-(V_1)$. Let $g \in C_0^\infty(B_\delta(x_0))$, $g \leq 0$. For $\epsilon > 0$, set $y_\epsilon(x) = x + \epsilon g e_n$ and $u_\epsilon(x) =$

$u_R(y_\epsilon(x))$. For ϵ sufficiently small, the monotonicity of u_R in the x_n -direction and the fact that $B_\delta(x_0) \subset C_R^-(V_1)$, imply that $V_1 \leq u_\epsilon \leq V_2$ on $B_\delta(x_0)$, and hence

$$0 \leq J(u_\epsilon, B_\delta(x_0)) - J(u_R, B_\delta(x_0)).$$

After rescaling appropriately and passing to the blow-up limit (using Lemma 2.5), the inequality above implies that

$$0 \leq J(U_\epsilon, B) - J(U, B),$$

where $U_\epsilon(x) = U(x + \epsilon \tilde{g} e_n)$, for any $\tilde{g}$ compactly supported in the unit ball B and $\tilde{g} \leq 0$. The same computations as in Lemma 3.11 give

$$(3.3) \quad 0 \leq \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} (J(U_\epsilon, B) - J(U, B)) = \int_{\partial\{U>0\}} (1 - a^2) \tilde{g} \nu_n$$

for all function $\tilde{g}$ as above. Thus, we contradict (3.2). $\square$

Combining the two Lemmas above, we obtain the following Corollary.

Corollary 3.12. *u_R is a variational solution to (2.1) in C_R .*

4. LOCAL REGULARITY THEORY

Throughout this section we will denote by u_R a viscosity solution to the problem

$$(4.1) \quad \begin{cases} \Delta u = 0 & \text{in } \Omega^+(u), \\ |\nabla u| = 1 & \text{on } F(u), \end{cases}$$

in the cylinder $C_R = \mathcal{B}_R(0) \times \{|x_n| < h_R\}$. Moreover, u_R is monotone increasing in $C_R^+(u_R)$ in the x_n direction, $V_1 \leq u_R \leq V_2$ in C_R , and $u_R = V_2$ on the sides $S_R = \partial\mathcal{B}_R(0) \times \{|x_n| \leq h_R\}$. Recall that V_1 is a strict smooth subsolution and V_2 is a strict smooth supersolution to (4.1) in $\mathbb{R}^n$, such that

- (i) $V_1 \leq V_2$ on $\mathbb{R}^n$, $0 \in \{V_2 > 0\} \cap \{V_1 = 0\}^\circ$;
- (ii) $\partial_n V_i > 0$ in $\overline{\{V_i > 0\}}$, for $i = 1, 2$.

The existence of u_R on a sufficiently tall cylinder is guaranteed by Theorem 3.1. Let u_R be extended to zero on $\{(x', x_n) : |x'| \leq R, x_n \leq -h_R\}$.

In this section we wish to prove the following regularity property for the free boundary of u_R .

Theorem 4.1. *For h_R sufficiently large, $F(u_R)$ is a Lipschitz graph in the x_n direction.*

Remark 1. The results in [C1] and [KN], imply that $F(u_R)$ is smooth in the interior.

Remark 2. The Lipschitz constant of $F(u_R)$ on C_R depends on the C^2 -norm of V_2 on $\overline{C_R}$, hence the result of Theorem 4.1 does not provide a “local” bound. Thus, we cannot control the Lipschitz constant on $F(u_R)$ in the limit as $R \rightarrow \infty$, using this bound. However, the Lipschitz constant is the same for all viscosity solutions u_R as in the beginning of this section.

We start by introducing a particular family of viscosity subsolutions (see [C1]) to the free boundary problem (4.1).

Lemma 4.2. *Let u be a viscosity solution to (4.1) in Ω . Let $v_t(x) = \sup_{B_t(x)} u(y)$, $t > 0$. Then v_t is a subsolution to (4.1) in its domain of definition. Furthermore, any point of $F(v_t)$ is regular from the positive side.*

We will also need the following results from [C1].

Lemma 4.3. *Let $v \leq u$ be two continuous functions in Ω , $v < u$ in $\Omega^+(v)$, v a subsolution and u a solution. Let $x_0 \in F(v) \cap F(u)$. Then x_0 cannot be a regular point for $F(v)$ from the positive side.*

Lemma 4.4. *(Comparison lemma) Let v_t be a family of subsolutions to (4.1), continuous in $\overline{\Omega} \times [0, T]$. Let u be a supersolution to (4.1) continuous in $\overline{\Omega}$. Assume that*

- (i) $v_0 \leq u$ in Ω ,
- (ii) $v_t \leq u$ on $\partial\Omega$ and $v_t < u$ in $\overline{\Omega^+(v_t)} \cap \partial\Omega$, for $0 \leq t \leq T$,
- (iii) every point $x_0 \in F(v_t)$ is regular from the positive side,
- (iv) the family $\Omega^+(v_t)$ is continuous.

Then, $v_t \leq u$ in Ω for every $t \in [0, T]$.

We now proceed to prove the following technical lemma. Here and henceforth the constant C will depend on the C^2 -norm of V_2 on $\overline{C_R}$.

Lemma 4.5. *(Existence of a barrier) There exists δ positive and small, such that for every $x_0 \in S_R \cap \overline{\{V_2 > 0\}} \cap \{|x_n| \leq h'_R\}$, $h'_R < h_R$, there exists a function V_{x_0} with the following properties*

- (a) $V_{x_0}(x_0) = u_R(x_0)$,
- (b) $u_R \geq V_{x_0}$
- (c) *for all $x \in S_R \cap \overline{\{V_2 > 0\}} \cap \{|x_n| \leq h'_R\}$, in a δ -neighborhood of x , the level set $V_x = u_R(x)$ is a Lipschitz graph in the vertical direction, with Lipschitz constant bounded by a uniform constant C independent of x .*

PROOF. Let $x_0 \in S_R \cap \overline{\{V_2 > 0\}} \cap \{|x_n| \leq h'_R\}$. Let H be the vertical half-space tangent to S_R at x_0 and containing C_R . Without loss of generality we can assume $H = \{x_1 > 0\}$. We wish to construct a function W_{x_0} such that $W_{x_0}^+$ is a subsolution and $V_2 \geq W_{x_0}^+$ on H , and also $V_2(x_0) = W_{x_0}(x_0)$. We have

$$(4.2) \quad V_2(x) \geq V_2(x_0) + l \cdot (x - x_0) - K|x - x_0|^2,$$

for some positive constant K , and vector l . Set

$$W_{x_0}(x) = (V_2(x_0) + (l - ae_1) \cdot (x - x_0) - K|x - x_0|^2)e^{-x_1}.$$

By choosing a sufficiently large, we can guarantee that $W_{x_0}^+$ is a strict subsolution in H , that is W_{x_0} is subharmonic in its positive phase and $|\nabla W_{x_0}| > 1$ on $F(W_{x_0})$. Indeed,

$$(4.3) \quad \Delta W_{x_0} = -2e^{-x_1}[l_1 - a - 2Kx_1 + nK] + W_{x_0},$$

hence on $H \cap \{W_{x_0} > 0\}$, if $a > l_1 + nK$ then $\Delta W_{x_0} > 0$. Also,

$$(4.4) \quad \partial_1 W_{x_0} = (l_1 - a - 2Kx_1)e^{-x_1} - W_{x_0},$$

hence on $F(W_{x_0})$, if $a > l_1 + 1$ then $\partial_1 W_{x_0} < -1$.

Let B be the (closed) ball on which W_{x_0} is non-negative, and let D be the $n-1$ dimensional ball $D := B \cap \partial H$. Let 0 be the center of D , and let us consider rescale of W_{x_0} around 0 , that is:

$$W_t(x) = tW_{x_0}\left(\frac{x}{t}\right).$$

One can easily verify that the family W_t^+ is non-increasing in t , in particular

$$W_t^+ \leq W_1^+ = W_{x_0}^+, \quad \text{for all } 0 \leq t \leq 1.$$

This follows from the fact that the function W_{x_0} is decreasing along each ray originating from x_0 into the half-space H . Hence, $V_2 \geq W_t^+$ in H for all $t \in [0, 1]$. Now, since $u = V_2$ on $S_R \subset H$, we can apply lemma 4.4 to u and the family W_t^+ , to conclude that $u \geq W_{x_0}^+$ in C_R , which gives the desired claim with $V_{x_0} = W_{x_0}^+$. $\square$

Theorem 4.1 is an immediate corollary to the following Theorem.

Theorem 4.6. *For h_R sufficiently tall, there exists a small constant $0 < c < 1$, such that, for small $s > 0$,*

$$(4.5) \quad \sup_{B_{cs}(x)} u_R(y - se_n) \leq u_R(x),$$

for all $x \in C_{R-cs}$.

PROOF. We divide the proof in 4 different steps.

Step 1. In this step we show that by taking h_R sufficiently tall, then (4.5) holds away from the free boundary. Indeed, let h_R be sufficiently large, so that a strip $T = \{|x'| \leq R, r_1 \leq x_n \leq r_2\} \Subset C_R^+(V_1)$, and $F(V_2)$ exits from the sides of C_R . Then,

$$u_R \geq \min_T V_1 = C \quad \text{on } T.$$

Now, let $(0, r) \in T$ and let $(0, y_n)$ belong to the level set $\{u_R = C/2\}$. Then,

$$(4.6) \quad C/2 \leq u_R(0, r) - u_R(0, y_n) = \int_{y_n}^r \partial_{x_n} u_R(0, t) dt.$$

Hence $\partial_{x_n} u_R(0, t) \geq C/r$, for some $t \in (y_n, r)$. Moreover, u_R grows linearly away from the free boundary, hence a ball centered at $(0, y_n)$ and of radius comparable to C is contained in the positive phase of u_R . Since u_R is monotone increasing in the vertical direction, we can cover the segment joining y_n and r with a finite (depending on $R, F(V_1), F(V_2)$) number of balls with radii comparable to C . Thus, since $\partial_n u_R$ is a positive harmonic function in $C_R^+(u_R)$, Harnack's inequality implies that $\partial_n u_R \geq c$ at $(0, r)$ for some constant c . Now Harnack's inequality up to the boundary implies that

$$(4.7) \quad \partial_n u_R \geq M \quad \text{on } T,$$

for some constant M . In particular, for r fixed between r_1 and r_2 , there exists a small c , such that

$$(4.8) \quad \sup_{B_{cs}(x)} u_R(y - se_n) < u_R(x),$$

for all $x \in \{x_n = r\} \cap \{|x'| \leq R - cs\}$, and s small.

Step 2. In this step we construct a family of subsolutions which we wish to compare to the solution u_R via a continuity argument, in the spirit of Lemma 4.4. To

guarantee that we can start the argument, we need to introduce the following quantity, which represents the length of the longest vertical segment contained in the free boundary of u_R :

$$\bar{s} = \sup\{\lambda > 0 \mid \exists \bar{x}, \text{ s.t. } (\bar{x} + \nu e_n) \in F(u_R), \forall |\nu| \leq \lambda\}.$$

Since $V_1 \leq u_R \leq V_2$, $0 \leq \bar{s} < +\infty$. Let s be a small positive number, and define $u_s(x) = u_R(x - (s + \bar{s})e_n)$. Now, consider the family of subsolutions

$$v_t^s(x) = \sup_{B_t(x)} u_s(y), \quad t \geq 0 \quad \text{and small.}$$

Since $u_R \leq V_2$ in C_R , by Lemma 4.5, we obtain that all the level sets of u_R in a small neighborhood of the boundary $S_R^+ = S_R \cap \{\overline{V_2 > 0}\} \cap \{x_n \leq r_2\}$, are trapped between two Lipschitz graphs, with Lip constants uniformly bounded. Hence, there exists a small constant c , such that if $\text{dist}(x, S_R) = cs$, then

$$(4.9) \quad \sup_{B_{\frac{c}{10}s}(x)} u_R(y - (s + \bar{s})e_n) \leq u_R(x).$$

Call $s' = cs/10$. Denote by

$$\Omega = \{-h_R < x_n < r\} \cap \{x \in C_R : \text{dist}(x, S_R) > cs\}.$$

Then, according to (4.8)-(4.9),

$$(4.10) \quad v_t^s(x) \leq u_R(x) \quad \text{for } x \in \partial\Omega, \quad \text{for all } t \in [0, s'].$$

Moreover,

$$(4.11) \quad v_t^s(x) < u_R(x) \quad \text{for } x \in \overline{\Omega^+(v_t^s)} \cap \partial\Omega.$$

Indeed (4.9) is strict on $\overline{\Omega^+(v_{s'})} \cap \partial\Omega$, since whenever $u_R(x)$ is zero, $B_{s'}(x - (s + \bar{s})e_n)$ is contained in the zero phase of V_2 , hence it cannot be tangent to $F(u_R)$. Also, the monotonicity of u_R in the x_n direction, guarantees that

$$(4.12) \quad v_0^s \leq u_R \quad \text{on } \Omega.$$

Step 3. In this step, we compare u_R with the family v_t^s , using a standard continuity argument. Define $A = \{t \in [0, s'] \mid v_t^s \leq u_R, \text{ in } \Omega\}$. By (4.12) $A \neq \emptyset$, and by the continuity in t of the family v_t^s , A is closed. We want to prove that A is open, hence $A = [0, s']$.

Let $t_0 \in A$, then $v_{t_0}^s \leq u_R$ in Ω , and by (4.11) and the strong maximum principle we get $v_{t_0}^s < u_R$ in $\Omega^+(v_{t_0}^s)$. If $t_0 > 0$, then by Lemma 4.2, every point of $F(v_{t_0}^s)$ is regular from the positive side, and Lemma 4.3 implies that $F(v_{t_0}^s) \cap F(u_R) = \emptyset$. Hence,

$$\overline{\Omega^+(v_{t_0}^s)} \subset \{x \in \overline{\Omega} \mid u_R(x) > 0\}.$$

The inclusion above, for the case $t_0 = 0$ follows from the definition of $\bar{s}$. By the continuity in t , for t close to t_0 ,

$$\overline{\Omega^+(v_t^s)} \subset \{x \in \overline{\Omega} \mid u_R(x) > 0\}.$$

Thus, $v_t^s < u_R$ on $\partial\Omega^+(v_t^s)$. Since $v_t^s - u_R$ achieves its maximum on the boundary, we then get $v_t^s < u_R$ on $\Omega^+(v_t^s)$, from which we conclude that $t \in A$.

Step 4. In this step we prove the desired statement (4.5). From Step 3 we have:

$$(4.13) \quad \sup_{B_{\frac{c}{10}s}(x)} u_R(y - (s + \bar{s})e_n) \leq u_R(x), \quad \text{in } \Omega.$$

Introduce the family of subsolutions

$$w_t(x) = \sup_{B_{c\frac{s}{10}}(x)} u_R(y - se_n + (t-1)\overline{se}_n).$$

Similar arguments as in Step 2, guarantee that hypothesis (ii) in Lemma 4.4 is satisfied. Moreover, according to (4.13), $w_t \leq u_R$ for $t = 0$, hence we can apply Lemma 4.4 to conclude that this inequality is true for $t = 1$ as well, which is the desired claim. $\square$

5. LOCAL NTA PROPERTY.

Let u_R be as in Theorem 3.1. In this section we prove the following property.

Theorem 5.1. *$F(u_R)$ is locally NTA.*

The proof of this result follows the line of [ACS], where NTA regularity for the free boundary of an optimization problem in heat conduction is proved.

We need the following monotonicity formula from [ACF].

Theorem 5.2. *Let v be a continuous function defined on $B = B_R(x_0)$. Suppose that v is harmonic in the open set $\{x \in B | v(x) \neq 0\}$. Let A_1 and A_2 be two different components in B of the set $\{x \in B | v(x) \neq 0\}$. Assume that for some constant $c > 0$, and any $r, 0 < r < R$,*

$$|B_r(x_0) \setminus (A_1 \cup A_2)| \geq c|B_r(x_0)|.$$

Define, for $0 < r < R$,

$$\phi(r) = \left(\frac{1}{r^2} \int_{B_r(x_0) \cap A_1} |\nabla v|^2 \rho^{2-n} dx \right) \left(\frac{1}{r^2} \int_{B_r(x_0) \cap A_2} |\nabla v|^2 \rho^{2-n} dx \right)$$

where $\rho = \rho(x) = |x - x_0|$. Then, for some positive β depending only on the dimension and the constant c , $r^{-\beta} \phi(r)$ is a non-decreasing function of r .

We also need the following result for harmonic functions, which can be obtained with an iterative argument in the spirit of Lemma 7 in [C3]. First we introduce a notation. For any real-valued function u defined on a domain $\Omega \subset \mathbb{R}^n$, and any $d \in \mathbb{R}$, we denote by

$$\Omega^d(u) = \{x \in \Omega | u(x) > d\}, \quad F^d(u) = \{x \in \Omega | u(x) \leq d\}.$$

Lemma 5.3. *Let u be a non-negative function in Ω , u harmonic in $\Omega^+(u)$, Lipschitz continuous and non-degenerate. Then, for any compact $D \subset \Omega$, there exist constants $\beta, \gamma > 0$ such that, whenever $\overline{x} \in F(u) \cap D^\circ$, $B_{2R}(\overline{x}) \subset D$, $x \in B_{R/2}(\overline{x}) \cap \{u > 0\}$ and A is a connected component of $\Omega^{\frac{u(x)}{2}}(u) \cap B_R(\overline{x})$ containing x , then there exists $B = B(\overline{y}, \gamma R) \subset A$, such that*

$$\int_B |\nabla u|^2 \geq \beta.$$

With the same notation as in Lemma 5.3, we have

Corollary 5.4. *There exists a constant τ , such that*

$$\int_A |\nabla u|^2 \rho^{2-n} dy > \tau R^2$$

where $\rho = \rho(y) = |y - \overline{x}|$.

PROOF OF LEMMA 5.3. Let $x = x_1$ and denote by $d_1 = |x_1 - x_0|$ the distance of x_1 from $F^{\frac{u(x_1)}{2}}(u)$. By the Lipschitz continuity and non-degeneracy of u , we have that the quantities,

$$d_1, u(x_1), d(x_1, F(u))$$

are comparable, with constants depending only on the Lipschitz and non-degeneracy constants of u on D . We wish to prove that there exists a positive constant δ and a point $x_2 \in B(x_1, d_1)$ such that

$$(5.1) \quad u(x_2) \geq (1 + \delta)u(x_1).$$

Indeed, by the Lipschitz continuity of u , there exists a constant c such that

$$u(y) \leq \frac{3u(x_1)}{4} \quad \text{on } B(x_0, cd_1).$$

Hence, the function

$$v := u - \frac{u(x_1)}{2}$$

is a non-negative harmonic function in $B(x_1, d_1)$, such that $v(x_1) = u(x_1)/2$ and $v \leq u(x_1)/4$ in $B(x_0, cd_1)$. Since,

$$v(x_1) = \oint_{B_r(x_1)} v, \quad r = d_1(1 - c/2),$$

we conclude that there exists $x_2 \in B_r(x_1)$ such that $v(x_2) \geq (1 + 2\delta)v(x_1)$ for a fixed positive constant δ . This, according to the definition of v , implies the desired claim (5.1). Moreover, from (5.1) we deduce that there exists y_1 on the line joining x_1 and x_2 , such that $|\nabla u|^2(y_1) > \beta$ for some small constant $\beta = \beta(\delta)$. Thus, since $|\nabla u|^2$ is subharmonic in $B(x_1, d_1)$, we get that

$$(5.2) \quad \oint_{B_{\delta_1}(y_1)} |\nabla u|^2 \geq \beta,$$

for $\delta_1 = cd_1/2$ comparable to d_1 , and $B(y_1, \delta_1) \subset B(x_1, d_1)$.

Now, we iterate this argument n -times, to find points x_j such that the following quantities are comparable,

$$(5.3) \quad d_j, u(x_j), d(x_j, F(u)),$$

and

$$(5.4) \quad u(x_{j+1}) \geq (1 + \delta)u(x_j), \quad \text{for all } j = 1, \dots, n.$$

Also,

$$(5.5) \quad \oint_{B_{\delta_j}(y_j)} |\nabla u|^2 \geq c, \quad B_{\delta_j}(y_j) \subset B(x_j, d_j), j = 1, \dots, n+1$$

with δ_j comparable to d_j . The number n is determined so that $B(x_{n+1}, d_{n+1})$ is the first ball to exit $B(x_0, R)$, but it is still included in $B(x_0, 2R)$. Thus, our lemma is proved with $\bar{y} = y_{n+1}$, if we show that d_{n+1} is comparable to R . From (5.4) and the fact that the quantities in (5.3) are comparable, we get that

$$cR \leq c|x_{n+1} - x_1| \leq c \sum_{i=1}^n d_i \leq \sum_{i=1}^n u(x_i) \leq Cu(x_{n+1}) \leq \tilde{C}d_{n+1}.$$

This concludes our proof. □

Finally, the proof of Theorem 5.1, is obtained combining the non-degeneracy and density property of u_R , together with the Harnack chain property from the next Lemma.

Lemma 5.5. *Let u be a viscosity solution to (2.1) in B_1 , such that u is Lipschitz continuous and nondegenerate, and u satisfies the density property (D). Then, there exists constants $M, \bar{\delta}$ such that, for any $\epsilon > 0$, and for any $x_1, x_2 \in B_{1/2}$, such that $B(x, \epsilon) \subset B_{3/4}^+(u)$ and $|x_1 - x_2| \leq C\epsilon \leq \bar{\delta}$, there exist $y_1 = x_1, \dots, y_l = x_2$, such that*

- (a) $B_i = B(y_i, \epsilon/M) \subset B_{3/4}^+(u)$, $i = 1, \dots, l$
- (b) $B_i \cap B_{i+1} \neq \emptyset$, $i = 1, \dots, l-1$
- (c) l independent of ϵ, x_1, x_2 .

PROOF. Assume that, without loss of generality,

$$\tilde{\delta} = \max\{d(x_1, \partial B_{1/2}^+(u)), d(x_2, \partial B_{1/2}^+(u))\} = d(x_2, \partial B_{1/2}^+(u)).$$

If $\tilde{\delta} \geq 2C\epsilon$, then, $x_1 \in B(x_2, C\epsilon) \subset B_{1/2}^+(u)$, and we can easily find the required chain.

Assume then, $\tilde{\delta} < 2C\epsilon$ and let $x_0 \in \partial B_{1/2}^+(u)$ be such that $\tilde{\delta} = |x_2 - x_0|$. Set $r_0 = 6C\epsilon$, then $x_1, x_2 \in B(x_0, r_0/2)$. Let $d = \frac{1}{2} \min\{u(x_1), u(x_2)\}$. We will show that, there exists $c \geq 1$, such that if $\tilde{\delta} \leq 1/(48c)$, and $R = cr_0 \leq 1/8$, then the connected components A_i of $B(x_0, R) \cap B_1^d(u)$ which contain $x_i, i = 1, 2$, are the same. Indeed, let us suppose that $A_1 \neq A_2$ and let us use Lemmas 5.2, and Corollary 5.4 with $v = (u - d)^+$. The density property of u guarantees that the hypotheses of Lemma 5.2 are satisfied, hence, for some exponent $\beta > 0$, the function $r^{-\beta}\phi(r)$ is non-decreasing. By Corollary 5.4 we obtain

$$\phi(r_0) > \tau^2.$$

Moreover, since u in Lipschitz on $B_{3/4}$ we also have the bound

$$\phi(R) \leq c',$$

with c' absolute constant independent of R . Hence,

$$\tau^2 r_0^{-\beta} < r_0^{-\beta} \phi(r_0) \leq R^{-\beta} \phi(R) \leq c' R^{-\beta}$$

or $R < c' r_0$, which is a contradiction if we choose $c = c'$.

We therefore conclude that $A_1 = A_2$. Since A_1 is open and connected we may find a curve Γ inside A_1 having x_1 and x_2 as end point. Denote by m the non-degeneracy constant of u on $B_{3/4}$. Then, for each $y \in \Gamma$ we know that

$$u(y) > d = \frac{1}{2} \min\{md(x_1, F(u)), md(x_2, F(u))\} \geq \frac{1}{2} m\epsilon.$$

Therefore, if K is the Lipschitz constant of u on $B_{3/4}$, for any $y \in \Gamma$, we have $d(y, F(u)) > \frac{1}{2} \frac{m\epsilon}{K}$. Set $\rho = \frac{1}{2} \frac{m\epsilon}{K}$, so that if $y \in \Gamma$ and $|x - y| < \rho$ then $u(x) > 0$. Since

$$\Gamma \subset \bigcup_{y \in \Gamma} B(y, \rho)$$

we may find a sequence $y_1, \dots, y_l$ of points in Γ such that $\Gamma \subset \bigcup_{i=1}^l B(y_i, \rho)$, and we may further ask that no y in Γ belong to more than $c(n)$ of the balls $B(y_i, \rho)$.

Furthermore, since $\rho = \frac{1}{2} \frac{m\epsilon}{K}$, $r_0 = 6C\epsilon$ and $y_i \in B(x_0, cr_0)$, l must be bounded by a constant depending only on dimension on c, C , but independent of x_1, x_2 or ϵ . $\square$

Remark 1. Notice that Lemma 5.5 shows that the free boundary of any viscosity solution to (2.1), which is Lipschitz continuous, non-degenerate, and satisfies the density property (D), is locally NTA.

Remark 2. It follows from the proof of Lemma 5.5 that M and $\bar{\delta}$ depend on the Lipschitz and non-degeneracy constants of u on $B_{3/4}$.

Finally, we recall a fundamental result about NTA domain (see [JK]).

Theorem 5.6. (*Dalberg Boundary Harnack principle*) *Let Ω be an NTA domain, and let V be an open set. For any compact set $G \subset V$, there exists a constant C such that for all positive harmonic functions u and v in Ω that vanish continuously on $\partial\Omega \cap V$, $u(x_0) = v(x_0)$ for some $x_0 \in \Omega \cap G$ implies $C^{-1}u(x) < v(x) < Cu(x)$ for all $x \in G \cap \bar{\Omega}$.*

6. GLOBAL MONOTONE SOLUTIONS.

In this section we prove our global result, that is Theorem 1.2. We start by deriving the following existence result, with a straightforward limit argument.

Theorem 6.1. *Assume that, there exist a strict smooth subsolution V_1 and a strict smooth supersolution V_2 to (2.1) in $\mathbb{R}^n$, such that*

- (i) $V_1 \leq V_2$ on $\mathbb{R}^n, 0 \in \{V_2 > 0\} \cap \{V_1 = 0\}^\circ$;
- (ii) $\partial_n V_i > 0$ in $\overline{\{V_i > 0\}}$, for $i = 1, 2$.

Then, there exists a global function u , viscosity solution to (2.1) in $\mathbb{R}^n$, such that u is monotone increasing in $\{u > 0\}$ in the x_n direction. Moreover $V_1 \leq u \leq V_2$, u is Lipschitz continuous, (I) non-degenerate, and it satisfies the density property (D).

PROOF. Let $\{R_k\}$ be a sequence of radii, $R_k \rightarrow +\infty$. Set $u_k := u_{R_k}$, where u_{R_k} is the viscosity solution on C_{R_k} from Theorem 3.1. Then, by Lemma 3.5, for any compact subset $D \subset \mathbb{R}^n$, and sufficiently large k , the functions $\{u_k\}$ are uniformly Lipschitz continuous on D . Hence, there exists a function $u : \mathbb{R}^n \rightarrow \mathbb{R}^+$, such that (up to a subsequence), $u_k \rightarrow u$ uniformly on compacts of $\mathbb{R}^n$, hence $V_1 \leq u \leq V_2$. Moreover, u is locally Lipschitz continuous, and monotone increasing in its positive phase in the x_n direction. Also, since the u_k 's are Lipschitz continuous, (I) non-degenerate, and satisfy the density property (D), with universal local constants, arguing as in Lemma 2.5, we obtain:

- (a) $\partial\{u_k > 0\} \rightarrow \partial\{u > 0\}$ in the Hausdorff distance;
- (b) $\chi_{\{u_k > 0\}} \rightarrow \chi_{\{u > 0\}}$ in L^1_{loc} ;
- (c) $\nabla u_k \rightarrow \nabla u$ a.e.

In particular, u is non-degenerate, (I) non-degenerate, and satisfies the density property (D). Furthermore, u is a variational solution to (2.1), on any compact, and it is harmonic in its positive phase. A blow-up argument as in Lemma 3.10 allows us to conclude that $F(u)$ cannot touch neither $F(V_1)$ nor $F(V_2)$. Hence, u is a viscosity solution to (2.1) in $\mathbb{R}^n$. $\square$

In particular, we can also conclude the following:

Corollary 6.2. *u minimizes $J(\cdot, B)$ among all competitors $v \in H^1(B)$, such that $V_1 \leq v \leq V_2$, and $v = u$ on ∂B , for all balls $B \subset \mathbb{R}^n$.*

To conclude the proof of Theorem 1.2, we need to prove the following:

Theorem 6.3. *$F(u)$ is a continuous graph, with a universal modulus of continuity on every compact $K \subset \mathbb{R}^n$.*

PROOF. We start by proving that $F(u)$ is a graph. Assume, by contradiction, that $F(u)$ contains a vertical segment.

Let $v(x) = u(x - te_n)$, for some small t . Since u is monotone in the x_n direction, we have $v \leq u$, and $v < u$ in $\{u > 0\}$. Moreover, by the assumption that $F(u)$ contains vertical segments, we have that $F(u) \cap F(v)$ is non-empty, for t sufficiently small. Assume, without loss of generality, that $0 \in F(u) \cap F(v)$. From Lemma 2.7, we obtain that u and v blow up around 0 to functions U and V which are homogeneous of degree 1. Also, U and V are viscosity solutions in $\mathbb{R}^n$, locally Lipschitz continuous, non-degenerate, and satisfying the density property (D). Hence, according to Remark 2 following Lemma 5.5, their free boundaries are NTA. Moreover, $U \geq V$. We wish to prove that,

$$(6.1) \quad U = \lambda V, \quad \text{on } \{V > 0\},$$

for some number $\lambda \geq 1$. Then, since U and V have the same asymptotic development near regular points, we get

$$(6.2) \quad U = V \quad \text{on } \{V > 0\}.$$

Towards proving (6.1), let us set

$$\lambda = \sup\{t > 0 \mid U \geq tV \text{ in } \{V > 0\}\}.$$

Clearly, since $U \geq V$, then $1 \leq \lambda < +\infty$. Define W to be the harmonic function in $B_1 \cap \{V > 0\}$ with boundary data

$$W = \begin{cases} U & \text{on } \partial B_1 \cap \{V > 0\}, \\ 0 & \text{on } F(V). \end{cases}$$

By the maximum principle,

$$\lambda V \leq W \leq U \quad \text{on } B_1 \cap \{V > 0\}.$$

If at some point $x_0 \in B_{1/2} \cap \{V > 0\}$, we have $(W - \lambda V)(x_0) = 0$, then $W \equiv \lambda V$ on $B_1 \cap \{V > 0\}$. Thus, since $W = U$ on $\partial B_1 \cap \{V > 0\}$, we have that $U = \lambda V$ at some point $x \in \partial B_1 \cap \{V > 0\}$. Therefore, since $U \geq \lambda V$ in $\{V > 0\}$, we conclude that $U \equiv \lambda V$ on $\{V > 0\}$.

If at some point $x_0 \in B_{1/2} \cap \{V > 0\}$, we have $W(x_0)/(\lambda V(x_0)) = \delta > 1$, then the boundary Harnack inequality (Theorem 5.6) implies that $W - \lambda V \geq c\delta\lambda V$ on $B_{1/2} \cap \{V > 0\}$. Therefore, since U and λV are homogeneous of degree 1, we conclude that $U \geq (1 + c\delta)\lambda V$ on $\{V > 0\}$, which contradicts the definition of λ .

Since $u > v$ in $\{v > 0\}$, a similar comparison argument for u and v allows us to conclude that

$$(6.3) \quad u \geq (1 + \epsilon)v \quad \text{on } \{v > 0\} \cap B_{1/2},$$

with ϵ depending on the ratio of u and v at a fixed scale. Passing to the blow-up limit, (6.3) produces a contradiction to (6.2).

Now, let us prove that $F(u)$ has a universal modulus of continuity on each compact cylinder K (box). We will denote by

$$v_K(\{x_n = \phi_1(x')\}, \{x_n = \phi_2(x')\}) := \max_{K'} |\phi_1(x') - \phi_2(x')|,$$

the vertical distance between two graphs in the x_n direction, with K' being the projection of the box K on the hyperplane $\{x_n = 0\}$.

We want to show that for every $K \subset \mathbb{R}^n$, and any $\epsilon > 0$, there exists a $\delta > 0$ such that, if $|\eta| < \delta$, then any u monotone global minimizer for J among competitors $v, V_1 \leq v \leq V_2$, satisfies,

$$v_K(\{u = \eta\}, F(u)) < \epsilon.$$

This, together with the non-degeneracy of u , gives the desired modulus of continuity estimate.

By contradiction, assume that for some $K \subset \mathbb{R}^n$, there exist a positive number ϵ , a sequence $\{\eta_j\}, \eta_j \rightarrow 0$, as $j \rightarrow +\infty$, and a sequence of energy minimizing solutions $\{u_j\}$, such that

$$(6.4) \quad u_j(x_j + \epsilon e_n) < \eta_j,$$

for some $x_j \in F(u_j) \cap K$.

The uniform Lipschitz continuity of the u_j 's, for j large, implies that (up to a subsequence):

$$u_j \rightarrow \tilde{u}, \quad \text{uniformly on compacts,}$$

and

$$x_j \rightarrow \bar{x} \in K,$$

with $\tilde{u}$ a Lipschitz continuous minimizing solution, monotone increasing in the x_n direction, and satisfying the (I) non-degeneracy, and the density property (D). Moreover $\tilde{u}(\bar{x}) = \tilde{u}(\bar{x} + \epsilon e_n) = 0$. We aim to prove that $\bar{x} \in F(\tilde{u})$; then by (6.4), we obtain that $F(\tilde{u})$ contains the vertical segment from $\bar{x}$ to $\bar{x} + \epsilon e_n$, which is a contradiction to what we showed above. Indeed, assume, that $\bar{x}$ does not belong to $F(\tilde{u})$. Then, there exists $r > 0$ such that,

$$(6.5) \quad B_r(\bar{x}) \subset \{\tilde{u} = 0\}^\circ.$$

If j is large enough, then $x_j \in B_{r/4}(\bar{x}) \cap F(u_j)$, and by non-degeneracy

$$\int_{B_{r/2}(\bar{x})} u_j \geq Kr.$$

Hence we get a contradiction to (6.5), as we pass to the limit for $j \rightarrow \infty$. $\square$

Acknowledgements. The results of this paper are part of my doctoral dissertation [D] at MIT. I would like to thank my advisor D. Jerison for introducing me to this subject, and for his guidance during my stay at MIT.

REFERENCES

- [AC] H.W. Alt, L.A. Caffarelli, *Existence and regularity for a minimum problem with free boundary*, J. Reine Angew. Math **325** (1981), 105–144.
- [ACF] H.W. Alt, L.A. Caffarelli, A. Friedman, *Variational problems with two phases and their free boundary*, Trans. Amer. Math. Soc. **282** (1984), no. 2, 431–461.
- [ACS] N.E. Aguilera, L.A. Caffarelli, J. Spruck, *An optimization problem in heat conduction*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **14** (1987), no. 3, 355–387.
- [AH] D.R. Adams, L.I. Hedberg, *Function spaces and potential theory*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **314** Springer-Verlag, Berlin, (1996)
- [BDG] E. Bombieri, E. De Giorgi, E. Giusti, *Minimal cones and the Bernstein problem*, Inv. Math. **7** (1969), 243–268.
- [C1] L.A. Caffarelli, *A Harnack inequality approach to the regularity of free boundaries. Part I: Lipschitz free boundaries are $C^{1,\alpha}$* , Rev. Mat. Iberoamericana **3** (1987) no. 2, 139–162.

- [C2] L.A. Caffarelli, *A Harnack inequality approach to the regularity of free boundaries. Part II: Flat free boundaries are Lipschitz*, Comm. Pure Appl. Math. **42** (1989), no.1, 55–78.
- [C3] L.A. Caffarelli, *A Harnack inequality approach to the regularity of free boundaries. Part III: Existence theory, compactness, and dependence on X* , Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **15** (1988), no. 4, 583–602 (1989).
- [CJK] L.A. Caffarelli, D. Jerison, C.E. Kenig, *Global energy minimizers for free boundary problems and full regularity in three dimension*, Noncompact problems at the intersection of geometry, analysis, and topology, 83–97, Contemp. Math., 350, Amer. Math. Soc., Providence, RI, 2004.
- [D] D. De Silva, *Existence and regularity of monotone solutions to free boundary problems*, Ph.D Thesis, 2005
- [DJ] D. De Silva, D. Jerison, *A singular energy minimizing free boundary*, Preprint (2005), submitted for publication.
- [DJ2] D. De Silva, D. Jerison, *Gradient bound for energy minimizing free boundary graphs*, in preparation.
- [F] A. Friedman, *Variational principles and free-boundary problems*, R. Krieger Publishing Co., Inc., Malabar, FL, 1988.
- [G] E. Giusti, *Minimal surfaces and functions of bounded variation*, Monographs in Mathematics, 80, Birkhäuser Verlag, Basel, 1984.
- [GT] D. Gilbarg, N. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der Mathematischen Wissenschaften, Vol. 224. Springer-Verlag, Berlin-New York, 1977.
- [JK] D. Jerison, C.E. Kenig, *Boundary behavior of harmonic functions in Non-tangentially Accessible Domains*, Adv. in Math. **46** (1982), no. 1, 80–147.
- [K] B. Kawohl, *Rearrangements and Convexity of Level Sets in PDE*, Lecture Notes in Mathematics, **1150**. Springer-Verlag, Berlin, 1985.
- [KN] D. Kinderlehrer, L. Nirenberg, *Analyticity at the boundary of solutions of nonlinear second-order parabolic equations*, Comm. Pure Appl. Math. **31** (1978), no. 3, 283–338.
- [W1] G.S. Weiss, *Partial regularity for weak solutions of an elliptic free boundary problem*, Commun. in Partial Diff. Equat. **23** (1998), no. 3-4, 439-455.
- [W2] G.S. Weiss, *Partial regularity for a minimum problem with free boundary*, J. Geom. Anal. **9** (1999), no. 2, 317–326.